

Flathead Family Planning
1035 1st Avenue West, Kalispell MT 59901
751-8150
Information for Depo-Provera

Depo-Provera is a progestin contraceptive injected intramuscularly (within a muscle) or subcutaneously (beneath the skin) every 11 to 13 weeks. About four in one hundred women may become pregnant while using Depo-Provera.

Women who use the Depo-Provera injection may have the following:

BENEFITS:

- * Less risk of endometrial cancer
- * Light or no menses (period), less anemia
- * Fewer menstrual cramps
- * Reduced risk of tubal pregnancy
- * Thirteen weeks of protection from one injection
- * Possible improvement of endometriosis
- * Fewer seizures for those with seizure disorders
- * No estrogen effects
- * Possible reduction in ovarian cancer
- * May be used by breastfeeding mothers

RISKS/SIDE EFFECTS (You may experience the following while using Depo-Provera):

- * Possible increase in depression or mood changes
- * Return to fertility may be delayed
- * Breast tenderness, acne, bloating, headache
- * Menstrual cycle disturbances
- * Change in appetite, weight gain or loss
- * Hair loss or increased hair growth
- * Side effects not immediately reversible
- * Decreased bone density
- * Rare allergic reaction
- * Dizziness, weakness or fatigue

You should **NOT** use Depo-Provera if you:

- * Have abnormal or unexplained bleeding from your vagina that has not been checked by a medical provider.
- * Have been taking *aminoluetethamide*, a drug for Cushing's syndrome.
- * Have a history of breast cancer, heart disease, stroke or blood clots.
- * Have current liver disease.
- * Think you may be pregnant.
- * Have an allergy to Depo-Provera

Use of Depo-Provera Contraceptive Injection may be considered among the risk factors for development of osteoporosis, (thinning of the bones). The rate of bone loss is greatest in the early years of use and then approaches the normal rate that would be expected for age.

ALTERNATIVES: You may receive information about other methods of birth control that are available. For situations of suspected contraceptive failure, emergency contraception is available and offers a second chance to avoid unintended pregnancy.

INSTRUCTIONS: You have received this information sheet about Depo-Provera and how it works. This injection does not protect you against STDs (sexually transmitted diseases) or HIV.

DECISION TO STOP USING: You may stop using Depo-Provera at any time. A woman is most likely to become pregnant if she or her partner does not use a method of birth control.

QUESTIONS: You may ask questions about Depo-Provera at any time and may contact the clinic with further questions.

INSTRUCTIONS FOR USING DEPO-PROVERA

1. There are four ways to start using Depo-Provera:
 - a. The first injection during the first five days of a normal menstrual period
 - b. Within the first five days postpartum if not breastfeeding
 - c. If exclusively breastfeeding at the sixth postpartum week
 - d. Your provider may provide Depo-Provera anytime in the cycle when known you are not pregnant. Use a backup method for seven days.
2. You may need to use a backup method of birth control for one week after your first injection. Your provider will instruct you if a backup method is necessary.
3. Because of the rare possibility of an allergic reaction, some providers ask that women remain in the clinic for 20 minutes after having their Depo-Provera injections.
4. Return to the clinic every 11 to 13 weeks for another injection. Mark your calendar for your next shot to be sure you are on time.
5. Be sure to use a backup method if you are late for injection.
6. It is important to exercise and take in adequate amounts of calcium while using Depo-Provera.

Late for an injection:

1. If you are more than one week late for your injection, use a backup method of contraception. Visit your clinic right away for your injection. You will need to continue using a backup method until you get your injection and for a week after that injection. Many clinicians will give you a pregnancy test to make sure you are not pregnant.
2. If you realize that you have missed your injection or are late for it, use another contraceptive or do not have intercourse. Return as soon as possible for your shot. Contact your clinic for possible emergency contraception.
3. If you have already had intercourse without being protected by a contraceptive, come to the clinic or call your clinician immediately to discuss options, including emergency contraception.

Depo-Provera and your Periods

1. Depo-Provera tends to make a woman's periods less regular, and spotting between periods is fairly common. Some women stop having periods completely. This is not harmful, and many women like not having periods.
2. If your pattern of bleeding is annoying, contact your clinician.
3. When you discontinue taking Depo, it may be a number of months before your periods return to normal.

Discontinuing Depo-Provera

1. If you discontinue Depo and do not want to become pregnant, start using a new contraceptive method 13 weeks from your last shot.
2. Depo injections may keep you from getting pregnant for more than 12 weeks after your last shot. The average delay in fertility is 10 months from the last injection. Depo does not decrease your fertility.

See your health-care provider if you develop any of the following symptoms:

- * Repeated, very painful headaches
- * Heavy bleeding
- * Depression
- * Severe, lower abdominal pain (maybe a sign of pregnancy)
- * Pus, prolonged pain or bleeding at injection site